

Left-right models with light neutrino mass prediction and dominant neutrinoless double beta decay rate

M. K. Parida* and Sudhanwa Patra†

Center of Excellence in Theoretical and Mathematical Sciences,
Siksha 'O' Anusandhan University, Bhubaneswar-751030, India

In TeV scale left-right symmetric models, new dominant predictions to neutrinoless double beta decay and light neutrino masses are in mutual contradiction because of large contribution to the latter through popular seesaw mechanisms. We show that in a class of left-right models with high-scale parity restoration, these results coexist without any contravention with neutrino oscillation data and the relevant formula for light neutrino masses is obtained via gauged inverse seesaw mechanism. The most dominant contribution to the double beta decay is shown to be via $W_L^- - W_R^-$ mediation involving both light and heavy neutrino exchanges, and the model predictions are found to discriminate whether the Dirac neutrino mass is of quark-lepton symmetric origin or without it. We also discuss associated lepton flavor violating decays.

I. INTRODUCTION: Evidences of tiny neutrino masses uncovered by the solar, atmospheric, and reactor neutrino oscillation experiments while calling for physics beyond the Standard Model (SM) might be strongly hinting at the fundamental nature of the particle i.e. whether Dirac [1] or Majorana [2]. In fact popular theories based upon seesaw mechanisms like type-I seesaw [3], type-II [4, 5], type-III [6, 7], inverse seesaw [8–11], and others [14–16, 18] come out with natural predictions of light Majorana neutrino masses. With lepton number violating mass insertion term by two units, confirmation of any events at the experimental search programmes for the neutrinoless double beta decay ($0\nu 2\beta$) would not only indicate the Majorana nature of the particles, but also it would strongly support the underlying seesaw mechanism for their mass generation. There have been attempts [19–23] on the experimental side to observe such a rare process, even with a present claim [24] while others are trying to improve the life time for this $0\nu 2\beta$ process [25–27]. So far, the Heidelberg-Moscow experiment using ^{76}Ge [24] has given the best limit on the half life, $T_{1/2} < 3 \times 10^{25}$ Yrs. which gives an upper bound on effective neutrino mass, $m_{\text{eff}} \leq 0.21 - 0.53$ eV. There are several interesting discussions and models using different seesaw mechanisms [28–31, 33–38] exploring possible non-standard contributions to $0\nu 2\beta$ transition.

Another important mysterious phenomenon of SM, namely, the origin of parity violation as monopoly of weak interactions, has been suggested to be having its underlying origin in the left-right (LR) symmetric interactions [39] which could be through the existence of mirror particles [40] of the SM or via left-right symmetric gauge theories [41, 42]. A very attractive aspect of LR gauge theory is its potential to explain the origin of parity (P) and CP violations in weak interactions and small neutrino masses. If left-right gauge theory has to make any significant impact on weak interactions phenomenology, the associated $W_R^\pm$ and Z_R boson masses have to be low. While current searches at the Large Hadron Collider restricts the lower bound on the scale of the RH gauge boson masses (M_R) to be $\mathcal{O}(1)$ TeV, $K_L - K_S$ mass difference gives $M_R > 2.5$ TeV [43]. Such a low scale W_R boson associated with right-handed charged currents can give additional contributions to $0\nu 2\beta$ and can be also accessible to LHC and future accelerator searches. As a result of this,

there can be various non-standard contributions to $0\nu 2\beta$ in LR gauge theories mediated by: (1.) two W_L gauge bosons (associated with left-handed currents), (2.) two W_R gauge bosons (associated with right-handed currents), (3.) one W_L and one W_R gauge boson at each vertex (mixed diagram) accompanied by both light and heavy neutrinos [33, 34]. In addition, there could be other contributions to $0\nu 2\beta$ in LR model due to doubly charged Higgs scalar exchanges where Majorana neutrino mass insertion has no role to play [32]. It is important to note here that the contributions to $0\nu 2\beta$ from the mixed diagram has been either ignored or considered to be sub-dominant [44], although this has been taken into account in the inverse process $e^- e^- \rightarrow W_L^- W_R^-$ in Ref. [45] for linear collider searches.

The natural TeV mass scale for RH Majorana neutrinos in conventional low scale LR gauge models emphasizing upon light neutrino mass generation mechanisms however predicts very large contribution to the light neutrino masses through canonical or type-II seesaw mechanisms [4, 41, 42]. Thus, it turns out that new dominant contributions to observable neutrinoless double beta decay ($0\nu 2\beta$) can not coexist with the experimentally determined tiny neutrino masses [46]. Alternatively, interesting proposals have been advanced where type-II seesaw dominance [33, 34] has been invoked by suppressing Dirac neutrino mass matrix in which case LR gauge theories may have only sub-dominant roles to play in representing charged fermion masses. The purpose of this letter is two fold: while showing that a crossed diagram with simultaneous W_L^- and W_R^- exchanges predicts the most dominant contribution to the ($0\nu 2\beta$), we provide a class of TeV scale left-right gauge theories where this is implemented without any suppression of naturally permitted Dirac neutrino masses and without any contravention with the neutrino oscillation data. The neutrino mass generation mechanism in these models turns out to be through gauged inverse seesaw.

II. THE MODEL: In conventional LR gauge theories, the type-I [3] and type-II seesaw [4] contributions to light neutrino masses are

$$m_\nu^I \sim -M_D \frac{1}{M_N} M_D^T, \quad m_\nu^{II} = f v_L$$

where the Dirac neutrino mass matrix M_D is similar to the

The relevant charged current interactions of leptons for this TeV scale LR gauge theory in the flavor basis is given by

$$\mathcal{L}_{CC} = \frac{g}{\sqrt{2}} \sum_{\alpha=e,\mu,\tau} \left[\bar{\ell}_{\alpha L} \gamma_{\mu} \nu_{\alpha L} W_L^{\mu} + \bar{\ell}_{\alpha R} \gamma_{\mu} N_{\alpha R} W_R^{\mu} \right] + \text{h.c.}$$

where, in terms of mass eigenstates $(\nu_{m_i}, S_{m_j}, N_{m_k})$ [35],

$$\begin{aligned} \nu_{\alpha L} &\sim \mathcal{N}_{\alpha i} \nu_{m_i} + \mathcal{U}_{\alpha j}^{\nu S} S_{m_j} + \mathcal{U}_{\alpha k}^{\nu N} N_{m_k}, \\ N_{\alpha R} &\sim \mathcal{V}_{\alpha i}^{N \nu} \nu_{m_i} + \mathcal{V}_{\alpha j}^{N S} S_{m_j} + \mathcal{R}_{\alpha k} N_{m_k}, \\ \mathcal{N}_{\alpha i} &= \left\{ \left(1 - \frac{1}{2} X X^{\dagger} \right) U_{\nu} \right\}_{\alpha i}, \\ \mathcal{R}_{\alpha k} &= \left\{ \left(1 - \frac{1}{2} X''^{\dagger} X'' \right) U_N \right\}_{\alpha k}, \\ \mathcal{U}_{\alpha j}^{\nu S} &= \{-X U_S\}_{\alpha j}, \quad \mathcal{U}_{\alpha k}^{\nu N} = \{-X' U_N\}_{\alpha k}, \\ \mathcal{V}_{\alpha i}^{N \nu} &= \{X^{\dagger} \frac{\mu_S}{M} U_{\nu}\}_{\alpha i}, \quad \mathcal{V}_{\alpha j}^{N S} = \{-X''^{\dagger} U_S\}_{\alpha j}. \end{aligned} \quad (7)$$

The non-unitarity matrices in our model are $X = \frac{M_D}{M}$, $X' = \frac{M_D}{M_N}$, and $X'' = \frac{M}{M_N}$ due to $\nu - S$, $\nu - N$, and $S - N$ mixings, respectively.

III. NEUTRINOLESS DOUBLE BETA DECAY: It is clear from the charged current interaction of this left-right gauge theory that, in addition to the standard contribution to $0\nu\beta\beta$ via light Majorana neutrino exchange, there are non-standard contributions due to the exchanges of heavy RH Majorana neutrinos and heavy sterile Majorana neutrinos. In addition, the extended seesaw ansatz manifests in non-standard contributions to lepton flavor violations and non-unitarity effects. In particular, we show that a dominant contribution to $0\nu\beta\beta$ arises due to mixed diagrams with simultaneous mediation of W_L^- and W_R^- bosons accompanied by light left-handed neutrinos and heavy right-handed Majorana neutrinos [34, 45] as shown in Fig. 1. We present analytic expressions for two most dominant contributions to the effective mass term and compare them with the standard contribution,

- m_{ν}^{ee} : which is analogous to the standard contributions, in this model,

$$m_{\nu}^{\text{ee}} = \mathcal{N}_{e i}^2 m_{\nu_i}, \quad (8)$$

- m_N^{ee} : which originates from the mediation of two W_R^- 's with the exchange of heavy RH Majorana neutrinos,

$$m_N^{\text{ee}} = p^2 \frac{M_{W_L}^4}{M_{W_R}^4} \frac{(\mathcal{R}_{e i})^2}{M_{N_i}}, \quad (9)$$

- $m_{\nu N}^{\text{ee}}$: which originates from simultaneous mediation of W_L^- and W_R^- and involves the Dirac mass matrix M_D

$$m_{\nu N}^{\text{ee}} \simeq p \left(\zeta_{LR} + \frac{M_{W_L}^2}{M_{W_R}^2} \right) \mathcal{N}_{e i} (M_N^{-1} M_D U_N)_{e i} \quad (10)$$

where, in our model, ζ_{LR} = LR mixing parameter $\leq 10^{-4}$.




FIG. 1: The neutrinoless double beta decay due to $W_L^- - W_R^-$ mediation in the mixed Feynman diagram with heavy and light neutrino exchanges described in the text. The cross in the right panel represents left-right mixing ζ_{LR} .

IV. RESULTS AND DISCUSSIONS: It is clear from equations (5) and (7) that the mass matrices M_D , M , and M_N are essential for predictions of light neutrino masses and $0\nu\beta\beta$. At first assuming the LR gauge theory to be having its high scale origin from Pati-Salam symmetry and neglecting the renormalization group corrections, the Dirac neutrino mass matrix is approximated as the up-type quark mass matrix via the CKM matrix and the running masses of the three up-type quarks, namely, $m_u = 2.33$ MeV, $m_c = 1.275$ GeV, and $m_t = 160$ GeV [48]

$$\begin{aligned} M_D &\sim V_{\text{CKM}} \hat{M}_u V_{\text{CKM}}^T \\ &= \begin{pmatrix} 0.067 - 0.004i & 0.302 - 0.022i & 0.55 - 0.53i \\ 0.302 - 0.022i & 1.48 - 0.0i & 6.534 - 0.0009i \\ 0.55 - 0.53i & 6.534 - 0.0009i & 159.72 + 0.0i \end{pmatrix} \text{ GeV} \end{aligned} \quad (11)$$

Under the condition $M_N \gg M \gg M_D$, the non-unitarity contribution of the extended seesaw model is mainly due to $\eta \simeq \frac{1}{2} X X^{\dagger}$, giving rise to $\eta_{\alpha\beta} = \frac{1}{2} \sum_{k=1}^3 \frac{M_{D\alpha k} M_{D\beta k}^*}{M_k^2}$ where we have assumed for the sake of simplicity: $M = \text{diag}[M_1, M_2, M_3]$. Then by saturating the available bound on $|\eta_{\tau\tau}| \leq 2.7 \times 10^{-3}$ [12, 49], we obtain

$$\frac{1}{2} \left[\frac{0.5805}{M_1^2} + \frac{42.72}{M_2^2} + \frac{25510.7}{M_3^2} \right] = 2.7 \times 10^{-3} \quad (12)$$

where the numbers inside the square bracket are in GeV^2 . We note that the above relation can be satisfied in the partial degenerate case, $M_1 = M_2 \geq 100$ GeV, and $M_3 \geq 2.2$ TeV and also in the non-degenerate case, $M_1 \geq 10$ GeV, $M_2 \geq 50$ GeV and $M_3 \geq 2.2$ TeV, but in the degenerate case, $M_1 = M_2 = M_3 = 2.2$ TeV.

a. Determination of μ_S from neutrino oscillation data: Inverting the neutrino mass formula given in eqn. (5) and using equation (7) and our model parameters, we obtain

$$\begin{aligned} \mu_S (\text{GeV}) &= X^{-1} \mathcal{N} \hat{m}_{\nu} \mathcal{N}^T (X^T)^{-1} \\ &= \begin{pmatrix} 0.01147 + 0.01i & -0.0027 - 0.0024i & 0.0007 + 0.002i \\ -0.0027 - 0.0024i & 0.0006 + 0.0005i & -0.0001 - 0.0004i \\ 0.0007 + 0.002i & -0.0001 - 0.0004i & -0.00004 + 0.0003i \end{pmatrix} \end{aligned}$$

where we have used the hierarchical neutrino masses $\hat{m}_{\nu}^{\text{diag}} = \text{diag}(0.00127 \text{ eV}, 0.00885 \text{ eV}, 0.0495 \text{ eV})$ and global fit to

the neutrino oscillation data including recent values of $\theta_{13} = 9.0^\circ$ and $\delta = 0.8\pi$ [46].

Thus, in the inverse seesaw approach, the light neutrino masses and large neutrino mixings including non-zero values of θ_{13} can be easily fitted through the elements of the μ_S matrix which may have interesting consequences on leptogenesis [13]. Although we have explicitly fitted the hierarchical light neutrino masses, similar fits can be obtained in the inverted hierarchical as well as the quasi-degenerate cases with corresponding elements of μ_S . In the case of M_D being similar to charged lepton mass matrix which holds true in conventional LR gauge theories [4, 42] neutrino oscillation data are similarly fitted with the corresponding μ_S matrix.

b. Neutrinoless double beta decay predictions: As explained in equations (7) -(10), the mixing matrices $X = \frac{M_D}{M}$, $X' = \frac{M_D}{M_N}$, and $X'' = \frac{M}{M_N}$ all contribute to non-standard predictions of $0\nu\beta\beta$ amplitude in the present left-right gauge theory. We have assumed the RH heavy Majorana neutrino mass matrix to be diagonal, $M_N = \text{diag}[M_{N_1}, M_{N_2}, M_{N_3}]$. Using the model parameters given in eqn. (11) for M_D , $M = \text{diag}[150, 150, 2500]$ GeV, $M_N = \text{diag}[5000, 5000, 10000]$ GeV, $U_\nu = U_{\text{PMNS}}$, $U_N = \mathbf{1}_{3 \times 3}$, and $U_S = \mathbf{1}_{3 \times 3}$, we have derived the relevant elements of the mixing matrices $\mathcal{N}_{ei}$, $\mathcal{R}_{ek}$, $\mathcal{U}_{ej}^{\nu S}$, $\mathcal{U}_{ek}^{\nu N}$, $\mathcal{V}_{ei}^{N\nu}$, and $\mathcal{V}_{ek}^{NS}$,

$$\begin{aligned} \mathcal{N}_{e1} &= 0.819, \quad \mathcal{N}_{e2} = 0.552, \quad \mathcal{N}_{e3} = 0.156, \\ \mathcal{R}_{e1} &= 0.997, \quad \mathcal{R}_{e2} = 0.0, \quad \mathcal{R}_{e3} = 0.0, \\ \mathcal{U}_{e1}^{\nu S} &= 0.00045, \quad \mathcal{U}_{e2}^{\nu S} = 0.002, \quad \mathcal{U}_{e3}^{\nu S} = 0.0002, \\ \mathcal{U}_{e1}^{\nu N} &= 0.00001, \quad \mathcal{U}_{e2}^{\nu N} = 0.00005, \quad \mathcal{U}_{e3}^{\nu N} = 0.000007, \\ |\mathcal{U}^{N\nu}| &\leq 10^{-9}, \quad \text{and} \quad |\mathcal{U}^{NS}| \leq 10^{-1}. \end{aligned} \quad (13)$$

With $|p| = 100$ MeV, $M_{W_R} = 5$ TeV and using equations (7) -(10), we predict the effective mass for $0\nu\beta\beta$ transition rate for hierarchical light neutrino masses,

$$\begin{aligned} |m_\nu^{\text{ee}}| &= \mathcal{N}_{e1}^2 m_{\nu_1} + \mathcal{N}_{e2}^2 m_{\nu_2} + \mathcal{N}_{e3}^2 m_{\nu_3} \\ &= 0.00157 \text{ eV}, \end{aligned} \quad (14)$$

$$|m_N^{\text{ee}}| = 6 \times 10^{-4} \text{ eV}, \quad (15)$$

$$|m_{\nu N}^{\text{ee}}| \sim 1 \text{ eV}. \quad (16)$$

Our numerical predictions are shown in Fig.2 as a function of W_R mass. With Dirac neutrino mass matrix having quark-lepton symmetric origin, the most dominant contribution due to $W_L^- - W_R^-$ mediation is found to be $m_{\nu N}^{\text{ee}} \simeq 1$ eV and 0.04 eV for $M_{W_R} = 5$ TeV, and 10 TeV, respectively. These predictions are reduced to $m_{\nu N}^{\text{ee}} \simeq 0.07$ eV and 0.03 eV for the corresponding values of the M_{W_R} when the Dirac neutrino mass matrix is similar to the charged lepton mass matrix. In other words, we predict that the $0\nu\beta\beta$ process would be able to discriminate LR gauge models having their roots in quark-lepton symmetry. We note that the sub-dominant contribution due to $W_R^- - W_R^-$ mediation given in eqn. (15) is suppressed as compared to the standard contribution due to $W_L^- - W_L^-$ mediation given in eqn. (14) for the same M_{W_R} masses shown in Fig. 3.




FIG. 2: Contributions to effective mass $m_{\nu N}^{\text{ee}}$ vs M_{W_R} using M_D similar to the up-quark mass matrix (upper curve) and M_D similar to charged lepton mass matrix (lower curve).




FIG. 3: Contributions to effective mass m_ν^{ee} and m_N^{ee} as a function of M_{W_R} . The slanting line represents our sub-dominant contribution due to $W_R - W_R$ mediation. For comparison, the standard contributions with hierarchical neutrino masses (the bottom horizontal line), inverted hierarchical masses (the middle horizontal line), and quasi-degenerate masses (the top horizontal line) are also given.

For the sake of comparison with the prediction for the inverse $0\nu\beta\beta$ processes in the golden channel $e^- e^- \rightarrow W_L^- W_R^-$ [45] which might be phenomenologically important for Linear Collider searches we used $M_{W_R} \geq 2.5$ TeV [43] to determine $\eta_\lambda = \frac{M_{W_L}^2}{M_{W_R}^2} \mathcal{N}_{ei} (M_N^{-1} M_D U_N)_{ei}$ which enters in cross-section for this process. Our model predicts this parameter to be 8.6×10^{-8} whereas the limit on this parameter is $\eta_\lambda \leq 9 \times 10^{-7}$ derived from the current experimental limit on $0\nu\beta\beta$ transition rate.

c. Lepton flavor violation: Besides the neutrinoless double beta decay process, the light and heavy neutrinos in this model can actively mediate different lepton flavor violating processes, $\mu \rightarrow e + \gamma$, $\tau \rightarrow e + \gamma$, and $\tau \rightarrow \mu + \gamma$ which are currently under active experimental investigation. The dominant contributions are mainly through the exchange of the six

heavy neutrinos [15] with branching ratio

$$\text{Br}(\ell_\alpha \rightarrow \ell_\beta + \gamma) = \frac{\alpha_w^3 s_w^2 m_{\ell_\alpha}^5}{256 \pi^2 M_W^4 \Gamma_\alpha} |\mathcal{G}_{\alpha\beta}^N + \mathcal{G}_{\alpha\beta}^S|^2 \quad (17)$$

$$\text{where } \mathcal{G}_{\alpha\beta}^N = \sum_k (\mathcal{U}^{\nu N})_{\alpha k} (\mathcal{U}^{\nu N})_{\beta k}^* \mathcal{F}\left(\frac{m_{N_k}^2}{M_{W_L}^2}\right),$$

$$\mathcal{G}_{\alpha\beta}^S = \sum_j (\mathcal{U}^{\nu S})_{\alpha j} (\mathcal{U}^{\nu S})_{\beta j}^* \mathcal{F}\left(\frac{m_{S_j}^2}{M_{W_L}^2}\right),$$

$$\mathcal{F}(x) = -\frac{2x^3 + 5x^2 - x}{4(1-x)^3} - \frac{3x^3 \ln x}{2(1-x)^4}.$$

Within the allowed range of model parameters $M_N \gg M \gg M_D$, it is clear that the first term in eq. (17) is negligible. The second term involving the heavy sterile neutrinos gives dominant contributions which is proportional to $\sum_j (\mathcal{U}^{\nu S})_{\alpha j} (\mathcal{U}^{\nu S})_{\beta j}^* \simeq 2\eta_{\alpha\beta}$ and our model predictions are

$$\begin{aligned} \text{Br}(\mu \rightarrow e + \gamma) &= 1.36 \times 10^{-15}, \\ \text{Br}(\tau \rightarrow e + \gamma) &= 1.06 \times 10^{-13}, \\ \text{Br}(\tau \rightarrow \mu + \gamma) &= 3.17 \times 10^{-12}. \end{aligned} \quad (18)$$

Noting that the present experimental limit at 90% C.L, $\text{Br}(\mu \rightarrow e + \gamma) \leq 1.2 \times 10^{-11}$ [50] is almost three orders of magnitude stronger than the limits $\text{Br}(\tau \rightarrow e + \gamma) \leq 3.3 \times 10^{-8}$ or $\text{Br}(\tau \rightarrow \mu + \gamma) \leq 4.4 \times 10^{-8}$ [51], appears to justify why the limit on $|\eta_{e\mu}|$ is at least one orders of magnitude better than the ones on $|\eta_{e\tau}|$ and $|\eta_{\mu\tau}|$. The projected reach of future sensitivities of ongoing searches are $\text{Br}(\tau \rightarrow e + \gamma) \leq 10^{-9}$, $\text{Br}(\tau \rightarrow \mu + \gamma) \leq 10^{-9}$, and $\text{Br}(\mu \rightarrow e + \gamma) \leq 10^{-18}$ [52, 53] throughout which the model predictions can be easily verified or falsified.

V. CONCLUSION: We have shown that in a class of left-right gauge theories, the light neutrino masses naturally arise though gauged inverse seesaw mechanism consistent with the current neutrino oscillation data. The associated TeV scale masses of $W_R^\pm$ and M_N can give dominant non-standard contributions to neutrinoless double beta decay which might be important for experimental searches. Specifically, we have demonstrated that the mixed diagram, via simultaneous mediation of W_L^- and W_R^- accompanied by the naturally predicted Dirac neutrino mass terms, gives the dominant contribution to $0\nu\beta\beta$ rate. Also this mixed diagram has rich phenomenological implication at ILC for the detection of the inverse process like $e^- e^- \rightarrow W_L^- W_R^-$. We have explicitly shown that this Dirac neutrino mass matrix could be similar to the up quark mass matrix which may have its high scale quark-lepton symmetric origin, or it may be similar to the charged lepton mass matrix expected from left-right gauge theory. The effective mass prediction in the former case being nearly 10 times larger than the latter case, we suggest that $0\nu\beta\beta$ signatures may probe high scale quark-lepton symmetry. As in our approach it is not necessary to fine tune the Dirac mass matrices, the left-right models could serve as promising theories for charged fermion masses. The TeV scale masses of $W_R^\pm$ and Z_R bosons are accessible to ongoing searches at LHC [54].

The predicted branching ratios for lepton flavor violating decays, being closer to the current experimental search limits, could be used to verify or falsify the left-right model framework considered in this letter.

* Electronic address: paridamk@soauniversity.ac.in

† Electronic address: sudha.astro@gmail.com

- [1] P. A. M. Dirac; Proceedings of the Royal Society of London, **109**, 752 (Dec. 1, 1925), pp. 642-653.
- [2] E. Majorana; N. Cim. **14** (1937) 171.
- [3] P. Minkowski, Phys. Lett. B **67**, 421 (1977); T. Yanagida, proceedings of the *Workshop on Unified Theories and Baryon Number in the Universe*, Tsukuba, 1979, eds. A. Sawada, A. Sugamoto, KEK Report No. 79-18, Tsukuba; S. Glashow, in *Quarks and Leptons, Cargèse 1979*, eds. M. Lévy. et al., (Plenum, 1980, New York); M. Gell-Mann, P. Ramond, R. Slansky, proceedings of the *Supergravity Stony Brook Workshop*, New York, 1979, eds. P. Van Nieuwenhuizen, D. Freeman (North-Holland, Amsterdam); R. N. Mohapatra and G. Senjanovic, Phys. Rev. Lett. **44**, 912 (1980).
- [4] R. N. Mohapatra and G. Senjanovic, Phys. Rev. D **23**, 165 (1981);
- [5] G. Lazarides, Q. Shafi and C. Wetterich, Nucl. Phys. B **181**, 287 (1981); J. Schechter and J. W. F. Valle, Phys. Rev. D **22**, 2227 (1980).
- [6] R. Foot, H. Lew, X. G. He and G. C. Joshi, Z. Phys. C **44**, 441 (1989); E. Ma and D. P. Roy, Nucl. Phys. B **644**, 290 (2002) [arXiv:hep-ph/0206150].
- [7] B. Bajc, G. Senjanović, JHEP **0708**, 014 (2007) [hep-ph/0612029]; B. Bajc, M. Nemevšek, G. Senjanović, Phys. Rev. **D76** (2007) 055011 [hep-ph/0703080]; A. Arhrib, B. Bajc, D. K. Ghosh, T. Han, G. -Y. Huang, I. Puljak, G. Senjanović, Phys. Rev. **D82**, 053004 (2010) [arXiv:0904.2390 [hep-ph]].
- [8] R. N. Mohapatra, Phys. Rev. Lett. **56**, 561-563 (1986); R. N. Mohapatra, J. W. F. Valle, Phys. Rev. **D34**, 1642 (1986).
- [9] D. Wyler, L. Wolfenstein, Nucl. Phys. **B218**, 205 (1983); E. Witten, Nucl. Phys. **B268**, 79 (1986).
- [10] J. L. Hewett, T. G. Rizzo, Phys. Rept. **183**, 193 (1989).
- [11] P. S. B. Dev, R. N. Mohapatra, Phys. Rev. **D81**, 013001 (2010) [arXiv:0910.3924 [hep-ph]].
- [12] Ram Lal Awasthi and Mina K. Parida, Phys.Rev. **D 86** (2012) 093004. e-Print: arXiv:1112.1826 [hep-ph].
- [13] S. Blanchet, P. S. B. Dev, R. N. Mohapatra, Phys. Rev. **D82**, 115025 (2010) [arXiv:1010.1471 [hep-ph]].
- [14] A. Pilaftsis, Z. Phys. C **55**, 275 (1992) [hep-ph/9901206]. J. Kersten and A. Y. Smirnov, Phys. Rev. D **76**, 073005 (2007) [arXiv:0705.3221 [hep-ph]]; R. Adhikari and A. Raychaudhuri, Phys. Rev. D **84**, 033002 (2011) [arXiv:1004.5111 [hep-ph]].
- [15] A. Ilakovac, A. Pilaftsis, Nucl. Phys. **B437**, 491 (1995) [hep-ph/9403398]; F. Deppisch, J. W. F. Valle, Phys. Rev. **D72**, 036001 (2005) [hep-ph/0406040]; C. Arina, F. Bazzocchi, N. Fornengo, J. C. Romao, J. W. F. Valle, Phys. Rev. Lett. **101**, 161802 (2008) [arXiv:0806.3225 [hep-ph]]; M. Malinsky, T. Ohlsson, Z. -z. Xing, H. Zhang, Phys. Lett. **B679**, 242-248 (2009) [arXiv:0905.2889 [hep-ph]]; M. Hirsch, T. Kernreiter, J. C. Romao, A. Villanova del Moral, JHEP **1001**, 103 (2010) [arXiv:0910.2435 [hep-ph]]; F. Deppisch, T. S. Kosmas, J. W. F. Valle, Nucl. Phys. **B752**, 80-92 (2006) [arXiv:0910.3924 [hep-ph]].

- [16] S. K. Kang, C. S. Kim, Phys. Lett. **B646**, 248-252 (2007) [hep-ph/0607072].
- [17] J. Ellis, J. L. Lopez and D. V. Nanopoulos, Phys. Lett. **B 292**, (1992) 189; J. Ellis, D. V. Nanopoulos and K. Olive, Phys. Lett. **B 300**, (1993) 121.
- [18] S. K. Majee, M. K. Parida, A. Raychaudhuri, Phys. Lett. **B668**, 299-302 (2008) [arXiv:0807.3959 [hep-ph]]; M. K. Parida, A. Raychaudhuri, Phys. Rev. **D82**, 093017 (2010) [arXiv:1007.5085 [hep-ph]].
- [19] H. V. Klapdor-Kleingrothaus, A. Dietz, L. Baudis, G. Heusser, I. V. Krivosheina, S. Kolb, B. Majorovits, H. Pas *et al.*, Eur. Phys. J. **A12** (2001) 147-154 [hep-ph/0103062].
- [20] C. E. Aalseth *et al.* [IGEX Collaboration], Phys. Rev. **D65** (2002) 092007 [hep-ex/0202026].
- [21] E. Conti *et al.* [EXO Collaboration], Phys. Rev. **B68**, 054201 (2003). [hep-ex/0303008].
- [22] C. Arnaboldi *et al.* [CUORE Collaboration], Nucl. Instrum. Meth. **A518**, 775-798 (2004) [hep-ex/0212053].
- [23] I. Abt, M. F. Altmann, A. Bakalyarov, I. Barabanov, C. Bauer, E. Bellotti, S. T. Belyaev, L. B. Bezrukov *et al.*, [hep-ex/0404039]; S. Schonert *et al.* [GERDA Collaboration], Nucl. Phys. Proc. Suppl. **145**, 242-245 (2005).
- [24] H. V. Klapdor-Kleingrothaus, I. V. Krivosheina, A. Dietz, O. Chkvorets, Phys. Lett. **B586**, 198-212 (2004) [hep-ph/0404088]; H. V. Klapdor-Kleingrothaus, I. V. Krivosheina, Mod. Phys. Lett. **A21**, 1547-1566 (2006).
- [25] C. Arnaboldi *et al.* [CUORICINO Collaboration], Phys. Rev. **C78**, 035502 (2008) [arXiv:0802.3439 [hep-ex]].
- [26] X. Sarazin *et al.* [NEMO Collaboration] [hep-ex/0006031]; J. Argyriades *et al.* [NEMO Collaboration], Phys. Rev. **C80**, 032501 (2009) [arXiv:0810.0248 [hep-ex]].
- [27] R. Arnold *et al.* [SuperNEMO Collaboration], Eur. Phys. J. **C70**, 927-943 (2010) [arXiv:1005.1241 [hep-ex]].
- [28] M. Hirsch, H. V. Klapdor-Kleingrothaus, S. G. Kovalenko, Phys. Lett. **B352**, 1-7 (1995) [hep-ph/9502315]; M. Hirsch, H. V. Klapdor-Kleingrothaus, S. G. Kovalenko, Phys. Rev. **D53**, 1329-1348 (1996) [hep-ph/9502385]; M. Hirsch, H. V. Klapdor-Kleingrothaus, S. G. Kovalenko, Phys. Rev. **D54**, 4207-4210 (1996) [hep-ph/9603213]; M. Hirsch, H. V. Klapdor-Kleingrothaus, S. G. Kovalenko, Phys. Rev. **D57**, 1947-1961 (1998) [hep-ph/9707207]; M. Hirsch, J. W. F. Valle, Nucl. Phys. **B557**, 60-78 (1999) [hep-ph/9812463].
- [29] B. C. Allanach, C. H. Kom, H. Pas, Phys. Rev. Lett. **103**, 091801 (2009) [arXiv:0902.4697 [hep-ph]].
- [30] V. Cirigliano, A. Kurylov, M. J. Ramsey-Musolf, P. Vogel, Phys. Rev. **D70**, 075007 (2004) [hep-ph/0404233]; V. Cirigliano, A. Kurylov, M. J. Ramsey-Musolf, P. Vogel, Phys. Rev. Lett. **93**, 231802 (2004) [hep-ph/0406199].
- [31] K. W. Choi, K. S. Jeong, W. Y. Song, Phys. Rev. **D66**, 093007 (2002) [hep-ph/0207180].
- [32] Noting that the doubly charged Higgs propagator provides additional damping factor to $0\nu\beta\beta$ amplitude, we ignore such contributions for sufficiently large values of the Higgs mass.
- [33] V. Tello, M. Nemevšek, F. Nesti, G. Senjanović, F. Vissani, Phys. Rev. Lett. **106**, 151801 (2011) [arXiv:1011.3522 [hep-ph]].
- [34] Joydeep Chakraborty, H. Zeen Devi, Srubabati Goswami and Sudhanwa Patra, JHEP 1208 (2012) 008. arXiv:1204.2527 [hep-ph].
- [35] W. Grimus and L. Lavoura, JHEP **0011**, 042 (2000), arXiv: 0008179 [hep-ph]; M. Hirsch, H. V. Klapdor-Kleingrothaus and O. Panella, Phys. Lett. B **374**, 7 (1996), arXiv: 9602306 [hep-ph]; M. Mitra, G. Senjanovic and F. Vissani, Nucl. Phys. B **856**, 26 (2012).
- [36] A. Ibarra, E. Molinaro and S. T. Petcov, Phys. Rev. D **84**, 013005 (2011).
- [37] A. Ibarra, E. Molinaro, S. T. Petcov, JHEP **1009**, 108 (2010) [arXiv:1007.2378 [hep-ph]]; A. Ibarra, E. Molinaro, S. T. Petcov, [arXiv:1101.5778 [hep-ph]].
- [38] M. Blennow, E. Fernandez-Martinez, J. Lopez-Pavon, J. Menendez, JHEP **1007** (2010) 096 [arXiv:1005.3240 [hep-ph]].
- [39] T. D. Lee and C. N. Yang, Phys. Rev. **D 104** (1956) 254.
- [40] R. Foot, R.R. Volkas, Phys. Rev. D 52, 6595 (1995). Z.G. Berezhiani, R.N. Mohapatra, Phys. Rev. D 52, 6607 (1995)
- [41] J. C. Pati and A. Salam, Phys. Rev. D 10 (1974) 275.
- [42] R. N. Mohapatra and J. C. Pati, Phys. Rev. D **11**, 2558 (1975); Phys. Rev. D **11**, 566 (1975); G. Senjanovic and R. N. Mohapatra, Phys. Rev. D **12**, 1502 (1975).
- [43] G. Beall, M. Bander, and A. Soni, Phys. Rev. Lett. **48** (1982) 848; A. Maiezza, M. Nemevsek, F. Nesti, and G. Senjanovic, Phys. Rev. **D 82**, 055022 (2010), arXiv: 1005.5160 [hep-ph]; D. Guadagnoli and R. N. Mohapatra, Phys. Lett. **B 694**, 386 (2011), arXiv: 1008.1074 [hep-ph]; S. Bertolini, J. O. Eeg, A. Maiezza and F. Nesti, Phys. Rev. **D 86** 095013 (2012), arXiv:1206.0668 [hep-ph].
- [44] M. Nemevsek, G. Senjanovic and V. Tello, arXiv: 1211.2837 [hep-ph].
- [45] James Barry, Luis Dorame and Werner Rodejohann; Eur. Phys. J. C72 (2012) 2023, arXiv:1203.3365 [hep-ph].
- [46] F. P. An *et al.* [DAYA-BAY Collaboration], arXiv:1203.1669 [hep-ex]; J. K. Ahn *et al.* [Soo-Bong Kim for RENO collaboration], arXiv:1204.0626v1 [hep-ex]; G. L. Fogli, E. Lisi, A. Marrone, A. Palazzo and A. M. Rotunno, Phys. Rev. D **84**, 053007 (2011), arXiv:1106.6028 [hep-ph]; D. V. Forero, M. Tortola and J. W. F. Valle, arXiv:1205.4018 [hep-ph].
- [47] D. Chang, R. N. Mohapatra, and M. K. Parida, Phys. Rev. Lett. **52**, 1072 (1984). D. Chang, R. N. Mohapatra, and M. K. Parida, Phys. Rev. **D 30**, 1052 (1984).
- [48] C. Amsler *et al.*, Phys. Lett. **B 667**, 1 (2008).
- [49] S. Antusch, J. P. Baumann, and E. Fernandez-Martinez, Nucl. Phys. B810, 369 (2009), 0807.1003.
- [50] M. L. Brooks *et al.* [MEGA Collaboration], Phys. Rev. Lett. **83**, 1521 (1999).
- [51] B. Aubert [The BABAR Collaboration], arXiv:0908.2381 [hep-ex].
- [52] For a review see F. R. Joaquim, A. Rossi, Nucl. Phys. B 765, 71 (2007).
- [53] Y. Kuno (PRIME Working Group), Nucl. Phys. B. Proc. Suppl. **149**, 376 (2005).
- [54] W. -Y. Keung and G. Senjanovic, Phys. Rev. Lett. **50**, 1427 (1983).